

BTM 2017 TRIP

















M018
07-030-S

**WELCOME
TAKULANDIRANI!**











H I J K L M N O P Q R S T U V W X Y Z 1 2 3 4 5 6 7 8 9 10 1































PNEUMONIA

CHI BAYO

Look for...

→ trouble breathing

Kupuma mowutina

→ nasal flaring

Kwakwetela hua mmphuno

→ can see ribs pulling in when breathing

holowa mwan kwa nthiti popuma

→ shortness of breath

qawumo

→ Cough, fever

C - t breathing

* If in doubt,
GO TO
HOSPITAL




Ngati mukukazi kira
pitani KU
chipitala
mwawsanga!

anga komoza na pipoofupi























PAEDIATRIC WARD

CHIPINDA CHA ANA

ST ANDREW'S HOSPITAL
CHILOOZA CHILOZA
MULABA - 000 000 - 1 00 000
MABALA - 000 000 - 1 00 000
MABALU - 000 000 - 1 00 000

PAEDIATRIC WARD

CHIPINDA CHA ANA

PUSH/KAMIKHANI





























NOT PUT FOOD
WATER IN
REFRIGERATOR











LIFO





F-75

F-75 Therapeutic milk

F-75 is especially formulated for the initial phase treatment of children with severe acute malnutrition.

F-75 is especially formulated for the initial feeding in the treatment of severe acute malnutrition. Never use F-75 for children to gain weight. It is not suitable for long term feeding of well-nourished children. Suitable for children aged 6 months and up. F-75 is only to be used under medical supervision of doctors and nutritionists.

Ready to feed

One sachet of F-75 with 500 ml boiled water provides 600 ml of F-75 therapeutic milk with an energy density of approx. 75 kcal/100ml.

How to prepare (see icons below)

1. Boil drinking water and let it cool to 70°C
2. Measure the quantity of water (500 ml / sachet 102.5g).
3. Add powder to cooled water (70°C) and stir vigorously (if possible with a whisk). Check the temperature before feeding. Use prepared F-75 therapeutic milk within 3 hours.



Important notice

- Breastfeeding is recommended for at least the first 24 months and exclusively until 6 months.
- Store powder dry at ambient temperatures below 30°C.
- Not intended for infants below 6 months of age.
- Discard remainders after 3 hours of preparation at room temperature.
- May be stored max. for 6 hours in the refrigerator.

102.5g net

F-75 Lait thérapeutique

F-75 est spécialement formulé pour le traitement de la phase initiale des enfants souffrant de malnutrition grave et sévère.

F-75 est spécialement formulé pour les premiers jours du traitement diététique de la malnutrition grave et sévère. Ne jamais utiliser F-75 dans le but de faire prendre du poids aux enfants. Ne convient pas à l'alimentation prolongée des enfants bien nourris. Convient aux enfants de plus de 6 mois. F-75 est à utiliser uniquement sous contrôle d'un médecin ou d'un nutritionniste.

Prêt à consommer

Un sachet de F-75 mélangé à 500ml d'eau fournit 600ml de lait thérapeutique d'une densité énergétique de 75 kcal/100ml.

Mode de préparation (voir schémas ci-dessous)

1. Faire bouillir de l'eau et laisser refroidir à 70°C.
2. Mesurer la quantité d'eau nécessaire (500ml pour un sachet 102.5g).
3. Ajouter la poudre à l'eau refroidie (70°C) et remuer vigoureusement (si possible avec un fouet). Vérifier la température avant consommation. La préparation F-75 reconstituée est à consommer dans les 3 heures.

Avis important

- L'allaitement maternel est recommandé durant les 24 premiers mois et exclusivement jusqu'à 6 mois.
- Stockez au sec à température ambiante inférieure à 30°C.
- Non destiné à des nourrissons de moins de 6 mois.
- Jeter la préparation après 3 heures à température ambiante.
- Peut être conservée pendant 6 heures maximum au réfrigérateur.



Valid Nutrition

PEANUT

RUTF

1 sachet = 500 kcal

Ready to Use Therapeutic Food

Aliment Thérapeutique Prêt-à-l'Emploi (ATPE)

For children >6 months with Severe Acute Malnutrition

Pour les enfants >6 mois atteints de Malnutrition Aiguë Sévère



ALL SAINTS CHURCH



























SHIPPER:
AVIMA (PTY) LTD
18 ASHCENBERG STREET
CHAMDOR, KRUGERSDORP
REP OF SOUTH AFRICA.
TEL: + 27 11 769 1300
CONTENTS:
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, SOLID, N.O.S.
(ALPHA CYPERMETHRIN)
UN 3077 CLASS 9 QTY = 6KG

CONSIGNEE:


















1:4

5

1:4
EXPERIENCED DRIVERS
FOR LEARNERS & BEGINNERS
PRACTICE

IN PLASTIC

DVD

intel

INVISIBLE










INTERNAL COMBUSTION ENGINE

























































































































































Strides
ARCTICAN LIMITED



Antimalarial Agent

COMBIART

Artemether / Lumefantrine Tablets

20/120mg

For Oral Use

1 x 24 Tablets

COMBIART
Artemether / Lumefantrine Tablets
20/120mg

"GOVERNMENT OF MALAWI"

Coartem® Dispersible

Artemether/Lumefantrine

20 mg/120 mg

Dispersible Tablets FOR CHILDREN

NOT FOR RETAIL SALE

NE PEUT ETRE VENDU DANS LE PRIVE

Weight: 5 kg to less than 15 kg

0 hrs Day 1 8 hrs Morning Day 2 Night Morning Day 3 Night

DAY 1 DAY 2 DAY 3

